

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ







## **Journal club**

GE11 peptide-conjugated nanoliposomes to enhance the  
combinational therapeutic efficacy of docetaxel and siRNA in  
laryngeal cancers

Prepared by: Dr. Alireza Farasat

25 February 2018

### Average Article Statistics

21 Days \*

from submission to first editorial decision.

36 Days \*

from editorial acceptance to publication.

\*Business days (Mon-Fri)

[Submit Manuscript](#)

### Journal Email Alerts

[Signup for Alerts](#)

[Back to Journals](#)

[PubMed](#)
[MEDLINE®](#)
[SJR OA](#)
[IMPACT FACTOR](#)

## International Journal of Nanomedicine

COPE  
 Member since 2011  
 JM07214



[About Journal](#)

[Editors](#)

[Peer Reviewers](#)

[Articles](#)

[Open Outlook: Nanomedicine](#)

[Publishing Fees](#)

[Aims and Scope](#)

[Call For Papers](#)

[Interview: Dr Webster](#)

[Award Nominations](#)

**ISSN: 1178-2013**

An international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. Reflecting the growing activity in this emerging specialty, the aim of this journal is to highlight research and development leading to potential clinical applications in the prevention and treatment of disease.

Endorsed by the [American Society for Nanomedicine](#)

Journal Impact Factor: 4.300 (5 year impact 5.008)

This journal is a member of and subscribes to the principles



Editor-in-Chief: [Dr Thomas J Webster](#)

[Scopus](#) Preview

[Author search](#)

[Sources](#)

[Help](#)

[Register](#)

[Login](#)

## Author details

[About Scopus Author Identifier](#)

The Scopus Author Identifier assigns a unique number to groups of documents written by the same author via an algorithm that matches authorship based on a certain criteria. If a document cannot be confidently matched with an author identifier, it is grouped separately. In this case, you may see more than one entry for the same author.

### Webster, Thomas Jay

Northeastern University, Department of Chemical Engineering, Boston, United States  
 Author ID: 7101765741

Other name formats: Webster, T. Webster, Thomas J. Webster, T. J. Webster, Thomas

Subject area:

[Materials Science](#)
[Engineering](#)
[Chemical Engineering](#)
[Biochemistry, Genetics and Molecular Biology](#)
[Chemistry](#)
[Pharmacology, Toxicology and Pharmaceutics](#)
[Physics and Astronomy](#)
[Medicine](#)
[Environmental Science](#)
[Social Sciences](#)

[View all](#)

Document and citation trends:

76



2052

[Follow this Author](#)

[View potential author matches](#)

**h-index:**

69

[View h-graph](#)

**Documents by author**

713

[Analyze author output](#)

**Total citations**

19874 by 11177 documents

[Print](#)
[Email](#)

IF= 4.3

## Editorial Board

40 Members

### Professor Abdollahi IRAN



Mohammad Abdollahi, Professor, Tehran University of Medical Sciences, Faculty of Pharmacy, and Pharmaceutical Sciences Research Center, Tehran, Iran.

### Dr Agrahari UNITED STATES



Vivek Agrahari, Ph.D., Formulation Research Scientist, Bayer HealthCare LLC, Shawnee, KS, USA.

### Prof. Dr. Bawa UNITED STATES



Raj Bawa, Bawa Biotechnology Consulting, Ashburn, Virginia, VA and Rensselaer Polytechnic Institute, Troy, NY, USA

# GE11 peptide-conjugated nanoliposomes to enhance the combinational therapeutic efficacy of docetaxel and siRNA in laryngeal cancers

This article was published in the following Dove Press journal:

International Journal of Nanomedicine

5 September 2017

Number of times this article has been viewed

Wei-Wei Xu<sup>1,\*</sup>

Da-yu Liu<sup>2,\*</sup>

Ying-chun Cao<sup>1</sup>

Xiang-yun Wang<sup>1</sup>

<sup>1</sup>Department of Ear-Nose-Throat,  
Dongying People's Hospital, Dongying,

<sup>2</sup>ENT & HN Surgery Department,  
Qilu Hospital, Shandong University,  
Jinan, Shandong, China

\*These authors contributed equally  
to this work

**Abstract:** In this study, dual therapeutic-loaded GE11 peptide-conjugated liposomes were developed and applied to enhance therapeutic efficacies of standard-of-care regimens for the treatment of laryngeal cancer. The therapeutic strategy used here was a combination treatment with the chemotherapeutic docetaxel (DTX) and siRNA against the ABCG2 gene that regulates multidrug resistance in many tumor types. Liposome-encapsulated DTX/ABCG2-siRNA molecules were targeted to recognize tumor cells of squamous morphology by conjugation to the EGFR-targeting ligand, GE11. Targeted, drug-infused liposomes were nanosized and exhibited controlled release of DTX. Presence of GE11 peptides on liposomal surfaces enhanced the quantities of liposomal constructs taken up by Hep-2 laryngeal cancer cells. GE11 peptide-conjugated liposomes also enhanced cytotoxic effects against Hep-2 laryngeal cancer cells when compared to treatment with free DTX, thereby reducing IC<sub>50</sub> values. Additionally, GE11 peptide-conjugated liposomes had significantly increased anti-tumor and apoptotic effects. Treatments with the GDLS nanoparticle formulation inhibited tumor growth in Hep-2 xenograft-bearing nude mouse models when compared to treatments with non-targeted NP constructs. Treatment of the mouse models with GE11 peptide-conjugated liposomes mitigated toxicities observed after treatment with free DTX. Taken together, liposomal encapsulation of DTX and ABCG2-siRNA improved the anti-tumor effects of treatment with free DTX in Hep-2 cell lines, and conjugation of GE11 peptides to liposomal constructs enhanced anti-tumor efficacies and specificities in laryngeal cancer cells.

**Keywords:** laryngeal cancer, docetaxel, ABCG2-siRNA, anti-tumor efficacy, liposomal chemotherapy carriers, GE11 peptide-targeted liposomes

# Background


- \* Mortality and incidence rate in laryngeal cancer.
- \* Treatment options for laryngeal cancers include surgery, radiotherapy, and chemotherapy.
- \* In chemotherapy, DTX acts by stabilizing the microtubule apparatus, thereby inhibiting cell division.
- \* Poor solubility and wide systemic side effects ,and multidrug resistance (MDR) are a barrier of chemotherapeutic agents that reduces therapeutic efficacy.

# Background

- \* ABCG2 gene
- \* The therapeutic strategy was used here was a combination treatment.
- \* Liposome as carrier
- \* EGFR and overexpression in laryngeal cancer
- \* A novel peptide, GE11 (YHWYGYTPQNVI) was specificity against EGFR proteins.




- \* In this paper, DTX loaded within the hydrophobic lipid bilayer.
- \* ABCG2–siRNA polynucleotides were loaded on liposomal surfaces using electrostatic interactions.
- \* Liposomal surfaces were also conjugated with GE11.






# Methods

1. Preparation of drug-loaded liposomes and peptide conjugations.
2. Particle size analyses.
3. Morphology analyses.
4. In vitro drug release assays.
5. In vitro cell viability assays of Hep-2 cells.
6. Apoptosis assays
7. Hoechst 33382 staining assays
8. In vivo anti-tumor efficacy studies
9. Statistical analyses



# Results and discussion


## 1. Physicochemical analyses of GE11 peptide-conjugated DTX/ABCG2–siRNA-loaded liposomes.



- The mean diameters of DTX/liposomes were ~110 nm
- Conjugations of GE11 peptides increased NP sizes to approximately 150 nm
- Final DTX/ABCG2–siRNA-loaded NPs were of mean size ~180 nm.

# Results and discussion


GE11-conjugated DTX/ABCG2-siRNA loaded liposomes were **slightly larger than DTX/liposomes** and maintained spherical morphologies, indicating that **surface modifications did not alter the shapes of the NPs.**






# Results and discussion

## 2- Drug-release studies



Dovepress



**Figure 3** In vitro drug release assays measuring loss of DTX from DL and GDSL constructs.


**Note:** The release study was performed using dialysis methods at 37°C.

**Abbreviations:** DL, DTX-loaded liposome; DTX, docetaxel.

➤ the in vitro drug release assays were determined by dialysis methods in PBS (pH 7.4) at 37°C

# Results and discussion


## 3- Cellular uptake analyses



➤ Rhodamine-B fluorescence intensities were observed in cytoplasmic regions, suggesting that NPs were internalized via endocytosis-mediated mechanisms.

# Results and discussion

## 4- In vitro cytotoxic effects of GE11-targeted DTX/ABCG2-liposomal NPs



**Figure 5** In vitro cytotoxic analyses of (A) empty NPs and (B) DTX/ABCG2-siRNA-loaded NPs.


**Notes:** The cells were treated with a concentration gradient of each NP formulation and then evaluated by MTT assays. \*\* $P < 0.001$  is the statistical difference between DL and GDSL.

**Abbreviations:** DL, DTX-loaded liposome; DTX, docetaxel; DSL, DTX and siRNA-loaded liposome; DTX, docetaxel; NP, nanoparticle.



# Results and discussion

## 5- Apoptosis assays



**Figure 6** Annexin-V/PI-based apoptosis assays of Hep-2 cells.

**Notes:** Cells were treated with free DTX, ABCG2-siRNA, DL, DSL, or GDSL constructs. The apoptosis rates were evaluated using flow cytometric analyses.

**Abbreviations:** DL, DTX-loaded liposome; DTX, docetaxel; DSL, DTX and siRNA-loaded liposome; FITC, fluorescein isothiocyanate; LL, lower left; LR, lower right; PI, propidium iodide; UL, upper left; UR, upper right.

# Results and discussion

## 6- Quantifications of nuclear apoptosis

- Nuclear apoptotic reactions of Hep-2 cancer cells were identified
- using Hoechst 33382 staining techniques.
- The control cell nuclei remained the same size and shape while nuclei treated with the NP formulations became irregular and distorted in shape.





Figure 7 Hoechst 33342-based apoptosis assays of Hep-2 cells.

**Notes:** Cells were treated with free DTX, ABCG2-siRNA, DL, DSL, or GDSL constructs. Nuclear morphologies of apoptotic cells were visualized using Hoechst stain  
**Abbreviations:** DL, DTX-loaded liposome; DSL, DTX and siRNA-loaded liposome; DTX, docetaxel.

# Results and discussion

## 7- In vivo anti-tumor efficacy assays



**Figure 8** In vivo anti-tumor efficacies of Hep-2 xenograft-implanted murine models.

**Notes:** Mice bearing Hep-2 tumors were treated with NP formulations, as listed, and **(A)** tumor volumes were measured on days 0–25. **(B)** Body weights were measured to determine toxicity of the treatments. \* $P < 0.05$ ; \*\*\* $P < 0.0001$ .

**Abbreviations:** DL DTX-loaded liposome; DSL DTX and siRNA-loaded liposome; DTX, docetaxel; NP, nanoparticle.



# Conclusion

1. The data presented here describe dual therapeutic-loaded GE11 peptide-conjugated liposomes that enhanced the combined therapeutic efficacies of DTX and ABCG2– siRNA in laryngeal cancers.
2. GE11 surface peptides enhanced cellular uptake of the constructs in Hep-2 laryngeal cancer cells.
3. Treatment of Hep-2 xenograft-implanted nude mice with the GDSSL NP formulation led to smaller tumors than those observed in animals receiving non-targeted NP constructs.

Thank you

